

Antioxidant properties of *Streptococcus thermophilus*, *Lactiplantibacillus plantarum* and *Levilactobacillus brevis* strains isolated from Brazilian tropical fruits

Giselle Santos Silva¹, Emma Faucher^{1,2}, Gabriela Zampieri Campos¹, Emília Maria França Lima¹, Katia Leani¹, Svetoslav Dimitrov Todorov^{1,3}

¹. ProBacLab, Universidade de São Paulo, São Paulo 05508-000, Brazil;

². AMU, Aix-Marseille University, Marseille 13006, France;

³. CISAS, Instituto Politécnico de Viana do Castelo, 4900-347 Viana do Castelo, Portugal;

Lactic acid bacteria (LAB) are renowned for their antioxidant properties, which play a crucial role in mitigating oxidative stress in the body. Oxidative stress arises from an imbalance between reactive oxygen species (ROS) and the body's ability to detoxify these reactive intermediates or repair the resulting damage. Excessive ROS can lead to cellular damage, contributing to various diseases, including cancer, cardiovascular diseases, and neurodegenerative disorders. LAB exhibit antioxidant activity through several mechanisms. They produce antioxidant enzymes such as superoxide dismutase (SOD) and can have pseudo-catalase activity, which neutralize ROS. Additionally, LAB can scavenge free radicals directly. For instance, strains like *Lactiplantibacillus plantarum* and *Limosilactobacillus fermentum* have demonstrated significant free radical scavenging abilities (Bryukhanov et al., 2022). Moreover, LAB can enhance the body's antioxidant defense by modulating the gut microbiota. They promote the growth of beneficial bacteria that produce short-chain fatty acids (SCFAs), which have anti-inflammatory and antioxidant effects. LAB also produce exopolysaccharides and other metabolites that contribute to their antioxidant capacity.

Streptococcus thermophilus L32, *Lactiplantibacillus plantarum* L3 and *Levilactobacillus brevis* L5 were isolated from Brazilian lemon, differentiated by repPCR and identified by biomolecular analysis, including partial sequencing of 16S rRNA. They were characterized regarding their antioxidant properties. Studied strains can be considered safe, based on the evaluation of hemolytic activity, degradation of mucin, susceptibility/resistance to antibiotics and production of the enzyme gelatinase. In addition, they can also be considered potentially beneficial based on their stability and survival in simulated gastrointestinal tract conditions (stomach and duodenum), α -galactosidase production, diacetyl production and specific levels of hydrophobicity.

Recorded levels of DPPH in the studied strains ranged between 93.52% for *Lpb. plantarum* L3, 77.88% for *Lvb. brevis* L5, to 53.29% for *Str. thermophilus* L32, indicating that antioxidant properties are strain specific, compared to the positive control, ascorbic acid (1 mg/mL) (74.80%). The beneficial attributes of these cultures to fermented food products may enable the reduction of chemical additives in line with consumers' demand for more natural and chemical-free food commodities.

Several studies have evaluated the DPPH antioxidant activity of *Str. thermophilus* with DPPH inhibition values ranging from 18.97% to 44.44%, indicating varying levels of antioxidant activity among the isolates (Kim et al., 2022). *Lpb. plantarum* showed significant DPPH radical scavenging activity, with some strains achieving scavenging percentages as high as 58.1%, or even comparable to those of ascorbic acid (Martiz et al., 2023). Research has shown that *Lvb. brevis* strains can achieve DPPH inhibition rates of up to 94.47%, making it one of the more effective LAB in terms of antioxidant potential (Won et al., 2021), while in different study was reported DPPH inhibition rates of 60.51% for *Lvb. brevis* (Qin et al., 2024).

The antioxidant properties of the studied strains can be applied to food preservation, reducing or even eliminating the need to use

The Yakult logo is displayed in a bold, red, sans-serif font.

International Symposia

Beneficial Microbes

March 27th and 28th, 2025

27 e 28 de Março, 2025

WYNDHAM SÃO PAULO IBIRAPUERA CONVENTION PLAZA

SÃO PAULO - BRAZIL

chemical additives, providing healthier options for consumers.

References

Bryukhanov, A.L. et al. Antioxidant properties of lactic acid bacteria. *Microbiology* 91, 463–478 (2022). <https://doi.org/10.1134/S0026261722601439>

Kim, S. et al. Antioxidant activity and probiotic properties of lactic acid bacteria. *Fermentation* 8(1), 29 (2022). <https://doi.org/10.3390/fermentation8010029>

Martiz, R.M. et al. Inhibition of carbohydrate hydrolyzing enzymes by a potential probiotic *Levilactobacillus brevis* RAMULAB isolated from fermented A

Won, G. et al. In vitro antidiabetic, antioxidant activity, and probiotic activities of *Lactiplantibacillus plantarum* and *Lactocaseibacillus paracasei* strains. *Curr. Microbiol.* 78, 3181–3191 (2021). h

Acknowledgements: FAPESP (2023/05394-9; 2023/14944-2); FCT, Portugal (UIDB/05937/2020 and UIDP/05937/2020); CAPES (33002010101P0)