

Protective Effect of *Weissella paramesenteroides* WpK4 on Murine Intestinal Mucositis: The Role of Tryptophan Metabolism

Gabriele Moreira Guimarães¹, Karen Costa¹, César da Silva Santana Moura^{1,2}, Sarah Elisa Diniz Moreira¹, Fernão Castro Braga¹, Marcelo Vidigal Caliar¹, Álvaro Cantini Nunes¹, Caio Tavares Fagundes¹, Elisabeth Neumann¹

¹ UFMG, Federal University of Minas Gerais, Minas Gerais, 31270-901, Belo Horizonte, Brazil;

² UFRJ, Federal University of Rio de Janeiro, Av. Athos da Silveira Ramos, CEP 21941909 Cidade Universitária - Rio de Janeiro;

The intestinal microbiota plays a crucial role in host homeostasis, with the balance between its components defining eubiosis, a state associated with host health. Dysbiosis, on the other hand, results from the loss of this balance and may be implicated in the development of various pathological conditions. Intestinal mucositis induced by chemotherapeutic agents, such as 5-fluorouracil (5-FU), is an example of an inflammatory condition in which dysbiosis can serve as an important biomarker. In this context, probiotics have been investigated as a therapeutic approach to restore eubiosis and mitigate the adverse effects of dysbiosis. Probiotics are live microorganisms that, when administered in adequate amounts, confer health benefits to the host, and evidence suggests they may be effective in controlling dysbiosis associated with inflammatory conditions. The bacterial strain *Weissella paramesenteroides* WpK4, isolated from the nasopharynx of weaned pigs, has shown probiotic potential, exhibiting beneficial effects in experimental models of infection and inflammation. A distinctive feature of this strain is the presence of the tryptophan operon in its genome, enabling it to synthesize tryptophan (TRP), an essential amino acid involved in critical biological processes, such as modulation of inflammatory responses and cellular signaling. Therefore, this study aimed to investigate the impact of tryptophan metabolism on the protection conferred by *W. paramesenteroides* WpK4 in a murine model of 5-FU-induced intestinal mucositis. Metabolic profiling of *W. paramesenteroides* WpK4 revealed the production of TRP, tryptamine (TAM), indole-3-acetic acid (IAA), and indole aldehyde (IAld), compounds known for their bioactive properties related to tryptophan metabolism. In the 5-FU-induced mucositis model, oral administration of *W. paramesenteroides* WpK4 resulted in 100% survival of the animals, while the control group, treated with 0.9% saline solution, showed a survival rate of 60%. Moreover, animals treated with *W. paramesenteroides* WpK4 exhibited lower morbidity and preservation of the intestinal microbiota composition, characteristics suggesting favorable modulation of the inflammatory response. Molecular analysis also indicated an increase in the expression of genes related to tryptophan metabolism in the host, suggesting the activation of the AhR (aryl hydrocarbon receptor) pathway, which may be involved in the observed protection. These results indicate that *W. paramesenteroides* WpK4 could represent a promising therapeutic strategy for the treatment of chemotherapeutic-induced intestinal mucositis, with its tryptophan metabolism possibly mediating the observed protective effects. The modulation of the intestinal microbiota and activation of tryptophan signaling pathways are mechanisms that warrant further investigation for the development of probiotic-based therapies in inflammatory contexts.

References

- ALVIM, L. B. et al. *Weissella paramesenteroides* WpK4 reduces gene expression of intestinal cytokines, and hepatic and splenic injuries in a murine model of typhoid fever. *Beneficial microbes*, v. 7, n. 1, p. 61-73, 2016.
- Sandes S, Figueiredo N, Pedrosa S et al (2020) *Weissella paramesenteroides* WpK4 plays an immunobiotic role in the gut-brain axis, reducing gut permeability, anxiety-like and depressive-like behaviors in murine models of colitis and chronic stress. *Food Res Biotechnol*.
- Rodrigues EP, Soares CDP, Galvão PG, et al (2016) Identificação de genes envolvidos na biossíntese do ácido indol-3-acético pela cepa *Gluconacetobacter diazotrophicus* PAL5 usando mutagenese de transposon. *Fr*.
- Giulietti A, Overbergh L, Valckx D et al (2001) Uma visão geral da PCR quantitativa em tempo real: aplicações para quantificar a expressão do gene da citocina. *Métodos* 25:386–401. <https://doi.org/10.1006/meth.2001>.

The Yakult logo is displayed in a bold, red, sans-serif font.

International Symposia

Beneficial Microbes

March 27th and 28th, 2025

27 e 28 de Março, 2025

WYNDHAM SÃO PAULO IBIRAPUERA CONVENTION PLAZA

SÃO PAULO - BRAZIL

Acknowledgements: We would like to express our sincere gratitude to CNPq and FAPEMIG for their essential support to research and innovation, which made the development of this work possible. Their encouragement is vital for the advancement of knowledge and for building a more promising future